


Today: Randomized Algorithms

Next class:

Quantum Algorithms

Wow! Cool! Neat!



The text 'Quantum Algorithms' is written in a mix of red and black ink. The word 'Quantum' is in red, and 'Algorithms' is in black. The 'Q' in 'Quantum' is a large, hand-drawn circle. The entire phrase is underlined with two wavy purple lines. There are four yellow stars: one to the left of 'Quantum', one above 'Quantum', one to the right of 'Algorithms', and one below 'Quantum'. Three green exclamations are scattered around: 'Wow!' to the left of the underline, 'Cool!' above the underline, and 'Neat!' below the underline.

Randomized Algorithms

Algorithms which uses random bits
to solve a problem

Allowed to fail with some probability
 $\leq 5\%$

Sometimes can be much faster
than deterministic

Integer factorization

Given a 500 digit number N
find its prime factorization $N = p_1 p_2 \dots p_k$

Algorithm: Check every integer $1 \leq x \leq \sqrt{N}$
to see if x divides N

Runtime: Suppose $N \approx 10^{500}$
Then $\sqrt{N} \approx \sqrt{10^{500}} = 10^{250}$
of atoms in universe $\leq 10^{106}$
 N has n digits $\Rightarrow$ needs $10^{n/2}$ time
Inefficient!

General number
field sieve:

Factor an n bit number
in time $\approx C n^{1/3} \log(n)^{2/3}$

Factoring is believed to be hard!

But very important!...

RSA-250: 250 digits (factored in 2020)

RSA-896: 270 digits \$75,000

RSA-2048: 618 digits \$200,000

(easy with a quantum computer)

Primality testing

Given an n digit number N
determine if it's prime or composite

Idea 1: Factor it! Runs in time $O(n^{1/3} \log(n)^{2/3})$ ☹️

Idea 2: Use something else about prime numbers...

Fermat's little theorem

If N is prime then $a^{N-1} \equiv 1 \pmod{N}$
for all $a \in \{1, \dots, N-1\}$

Fermat Test (N)

1. Pick $a \in \{1, \dots, N-1\}$ uniformly at random

2. If $a^{N-1} \equiv 1 \pmod{N}$, output "prime"

Otherwise, output "composite"

Fact: If N is prime, always outputs "prime"

$N = 12 = 3 \cdot 4$ ^{not coprime} $a = 2$ $2^{11} \not\equiv 1 \pmod{12}$

^{not coprime}

$a = 3$

$3^{11} \not\equiv 1 \pmod{12}$

^{coprime}

$a = 5$

$5^{11} \equiv 1 \pmod{12} ???$

$N = p^2$, p large prime $\Rightarrow$ only $\frac{1}{p}$ of a 's are not coprime

Test only good if $a^{N-1} \equiv 1 \pmod{N}$ ^{with N} for lots of coprime a

Carmichael numbers

Composite numbers N s.t.

$$a^{N-1} \equiv 1 \pmod{N}$$

for all a coprime to N

Pass the Fermat test for all coprime a

$$N = 561 = 3 \cdot 11 \cdot 17$$

Let's pretend these don't exist for now..


Thm: Suppose N is composite and not Carmichael.
Then $\Pr[\text{Fermat Test}(N) = \text{composite}] \geq 1/2$

Pf: Not Carmichael $\Rightarrow$ coprime b s.t. $b^{N-1} \not\equiv 1 \pmod{N}$

Claim: Suppose a passes Fermat Test: ($a^{N-1} \equiv 1 \pmod{N}$)
Then $a \cdot b \pmod{N}$ fails Fermat Test.

Pf:

$$\begin{aligned}(a \cdot b)^{N-1} &\equiv a^{N-1} \cdot b^{N-1} \pmod{N} \\ &\equiv b^{N-1} \pmod{N} \\ &\not\equiv 1 \pmod{N}\end{aligned}$$



a's pass test
 $a^{N-1} \equiv 1 \pmod{N}$

a's fail test
 $a^{N-1} \not\equiv 1 \pmod{N}$

Need to check

$$a_i \cdot b \not\equiv a_j \cdot b \pmod{N} \quad (i \neq j)$$

Fact: b is coprime
 $\Rightarrow$ inverse $b^{-1} \pmod{N}$

$$a_i \cdot \underset{\cdot b^{-1}}{b} \not\equiv a_j \cdot \underset{\cdot b^{-1}}{b} \pmod{N}$$

$$a_i \not\equiv a_j \pmod{N}$$

$$\text{So } |\text{pass}| \leq |\text{fail}|$$

□

$\Pr[\text{Fermat Test}(N) = \text{composite}] \geq \frac{1}{2}$ (if N is composite)

Repeat k times $\rightarrow$ detect composite
w/prob $\geq 1 - \frac{1}{2^k}$

Can be very confident!

Runtime

Need to compute $a^{N-1} \pmod{N}$

Suppose $N-1 = 2^n$

$$\left. \begin{array}{l} a \cdot a \equiv a^2 \pmod{N} \\ a^2 \cdot a^2 \equiv a^4 \pmod{N} \\ a^4 \cdot a^4 \equiv a^8 \pmod{N} \\ \vdots \\ a^{2^{n-1}} \cdot a^{2^{n-1}} \equiv a^{2^n} \pmod{N} \end{array} \right\} n \text{ steps}$$

(Can generalize to arbitrary $N-1$)

Can add another check to detect Carmichael numbers

This gives Miller-Rabin primality test (1976)

Since then, we've only had randomized algs (no deterministic)

undergrads *Until...*

Agrawal-Kayal-Saxena Primality test (2002)

A deterministic alg for primality testing
in $O(n^{12})$ time (later $O(n^6)$ time)

Miller 1976 "derandomization"

Try the Miller-Rabin test for all $a \leq O(n^2)$

This will detect if N is prime or composite

(assuming generalized Riemann hypothesis)

Primarily: efficient randomized algorithm first,
later efficient deterministic alg

Other problems: Only know efficient randomized alg
(Polynomial identity testing)

Two possible worlds: 1. Every efficient randomized alg
has deterministic counterpart

efficient
deterministic

→ P

2. Some problems only have
efficient randomized algs


v.

BPP

← efficient
randomized

Minimum cut

Unweight, undirected
graphs $G = (V, E)$



cut of
size 1

cut of size 3

Idea: Max flow/min cut alg

Computes min s - t cut in time $O(n \cdot m)$

Which s, t to use?

$|V|$ $|E|$

set $s=1$, try all $t=2, \dots, n$


$O(n^2 \cdot m)$ time

Karger's algorithm(G)


for $i = 1, \dots, n-2$ ($n = |V|$)

1. pick a uniformly random edge e
2. contract e

return cut specified by the remaining ^{two} supervertices



Intuition



Karger's alg finds min cut if it never contracts e
But way more edges on left
Will usually pick there!

Thm: Let $C = (S, \bar{S})$ be a min cut of size k .
 $\Pr[\text{Karger's alg outputs } C] \geq \frac{1}{\binom{n}{2}} = \frac{2}{n(n-1)}$

Pf: Let G_i for graph at beginning of i^{th} iteration
($G_1 = G$)

Fact 1: Min-Cut in $G_i \geq k$.
(any cut corresponds to cut in G)

Fact 2: # of vertices in $G_i = n - (i-1)$
 $= n - i + 1$

Fact 3 : degree of each vertex in $G_i \geq k$

Fact 4 : # of edges in G_i

$$= \frac{1}{2} \sum_{v \in G_i} d_v \geq \frac{1}{2} \sum_{v \in G_i} k$$

$$= \frac{1}{2} k \cdot |G_i|$$

$$= \frac{1}{2} k \cdot (n - i + 1)$$

Suppose at G_i , haven't contracted edge in C yet.

$$\Pr[\text{don't contract an edge in } C]$$

$$= 1 - \Pr[\text{contract an edge in } C]$$

$$\geq 1 - \frac{k}{\frac{1}{2} \cdot k \cdot (n-i+1)} = 1 - \frac{2}{(n-i+1)}$$

$$\Pr[\text{never contract edge in } C]$$

$$\geq \left(1 - \frac{2}{n}\right) \left(1 - \frac{2}{n-1}\right) \left(1 - \frac{2}{n-2}\right) \cdots \left(1 - \frac{2}{3}\right)$$

$$= \left(\frac{\cancel{n-2}}{n}\right) \left(\frac{\cancel{n-3}}{n-1}\right) \left(\frac{\cancel{n-4}}{n-2}\right) \cdots \left(\frac{2}{\cancel{4}}\right) \left(\frac{1}{\cancel{3}}\right)$$

$$= \frac{2}{n(n-1)}$$

$$\Pr[\text{succeeds}] \geq \frac{1}{\binom{n}{2}} \approx \frac{2}{n^2}$$

succeed w/ constant prob: repeat n^2 times
(or slightly more)

$$\# \text{ of min cuts} \leq \binom{n}{2}$$

$$\Pr(\text{output min cut}) \geq \sum_{\text{min cut}} \frac{1}{\binom{n}{2}} =$$