

Disc 3A — "Slow Paced"

Announcements

- 16A Review Session — 5-6:30PM today
 - Self-grades and resubmissions due tomorrow!
 - HW 3 released, due this Friday
 - Lite Lab 1 due today
 - Hit me up anytime at eeecs16b-sp21@berkeley.edu
-

1. Differential equations with piecewise constant inputs

Working through this question will help you understand better differential equations with inputs. Along the way, we will also touch a bit on going from continuous-time into a discrete-time view. This problem also provides a vehicle to review relevant concepts from calculus.

(a) Consider the scalar system

$$\frac{d}{dt}x(t) = \lambda x(t) + u(t) \quad \text{Input}$$

Rate of change Scaled signal


Goal: model the change in signal $x(t)$ as affected by input $u(t)$

Our goal is to solve this system (find an appropriate function $x(t)$) for general inputs $u(t)$. To do this, we will start with a piecewise constant $u(t)$; we already have the tools to solve this system, which we will do in the first few parts. Later in the worksheet, we will extend this to general $u(t)$.

Suppose that $x(t)$ is continuous (in real systems, this is almost always true). Further suppose that our input $u(t)$ of interest is piecewise constant over durations of width Δ . In other words:

$$u(t) = u(i\Delta) = u[i] \text{ if } t \in [i\Delta, (i+1)\Delta). \quad (2)$$

In keeping with this notation, we will use the notation



$$x_d[i] = x(i\Delta).$$

Example for intuition: think of measuring the voltage $x(t)$ across some circuit element at time t , while also administering some input voltage $u(t)$ at time t .




Figure 1: An example of a discrete input where the limit as the time-step Δ goes to 0 approaches a continuous function. The red line, the original signal $u_c(t) = \sin(t)$, is traced almost exactly by the blue line, which has a small time-step, and not nearly as well by the green line, which has a large time-step.

Here is a solution to this system, which may help with visual intuition:




Figure 2: An example of a solution to this diff. eq. system. In this case $\lambda = 1$, $u[0] = 10$, $u[1] = -30$.

a)

The first step to analyzing this system is to discover its behavior across a time-step with constant input, since we already know how to solve these kinds of systems.

Given that we know the value of $x(i\Delta) = x_d[i]$, compute $x_d[i+1] = x((i+1)\Delta)$.


Hint: For $t \in [i\Delta, (i+1)\Delta]$, the system is

$$\frac{d}{dt}x(t) = \lambda x(t) + u[i].$$

$$\frac{d}{dt}x(t) = \lambda x(t) + u[i]$$

↳ Given we know $x(i\Delta)$,

$$x((i+1)\Delta) = \underbrace{x(i\Delta)}_{\text{initial value}} + \int_{t=i\Delta}^{t=(i+1)\Delta} \underbrace{\frac{d}{dt}x(t)}_{\text{overall change}} dt$$



Notation

$$x(t = i \cdot \Delta) = x_d[i]$$

$$u(1\Delta) = u[i] = u_d[i]$$

Differential equation for time $t \in [i\Delta, (i+1)\Delta]$:

$$\frac{d}{dt}x(t) = \lambda x(t) + u(t) = \lambda x(t) + u[i]$$

⇒ Want to fit to solution $x(t) = \alpha \cdot e^{\lambda(t-i\Delta)} + \beta$

$$\frac{d}{dt}x(t) = \lambda \cdot \alpha \cdot e^{\lambda(t-i\Delta)} = \lambda x(t) + u[i]$$

$$\Rightarrow \cancel{\lambda \cdot \alpha} e^{\lambda(t-i\Delta)} = \lambda (\alpha e^{\lambda(t-i\Delta)} + \beta) + u[i]$$

$$0 = \lambda \beta + u[i] \Rightarrow \boxed{\beta = -\frac{u[i]}{\lambda}}$$

Now, use initial condition:

$$x(i\Delta) = \alpha e^{\lambda(i\Delta - i\Delta)} + \beta$$

$$x(i\Delta) = \alpha + \beta \Rightarrow \alpha = x(i\Delta) - \beta$$

$$\boxed{\alpha = x(i\Delta) + \frac{u[i]}{\lambda}}$$

$$x(t) = \alpha \cdot e^{\lambda(t-i\Delta)} + \beta$$

$$x(t) = \left(x(i\Delta) + \frac{u[i]}{\lambda}\right) \cdot e^{\lambda(t-i\Delta)} - \frac{u[i]}{\lambda}$$

$$= x(i\Delta) \cdot e^{\lambda(t-i\Delta)} + \frac{u[i]}{\lambda} \cdot e^{\lambda(t-i\Delta)} - \frac{u[i]}{\lambda}$$

$$x(t = (i+1)\Delta) = x(i\Delta) \cdot e^{\lambda((i+1)\Delta - i\Delta)} + u[i] \cdot \frac{1}{\lambda} (e^{\lambda((i+1)\Delta - i\Delta)} - 1)$$

$$x_d[i+1] = x_d[i] \cdot e^{\lambda \Delta} + u[i] \cdot \frac{1}{\lambda} (e^{\lambda \Delta} - 1)$$

$$\Rightarrow \boxed{x_d[i+1] = x_d[i] \cdot e^{\lambda \Delta} + \frac{u[i]}{\lambda} \cdot (e^{\lambda \Delta} - 1)}$$

From Note 2, pg 1 eq 1

Dif eq of form	$\frac{d}{dx}x(t) = \lambda x(t) + u(t)$
has soln	$x(t) = \alpha \cdot e^{\lambda(t-i\Delta)} + \beta$
α and β are fns of $x(i\Delta), u[i], \lambda$	

- (b) Now that we've found a one-step recurrence for $x_d[i+1]$ in terms of $x_d[i]$, we want to get an expression for $x_d[i]$ in terms of the original value $x(0) = x_d[0]$, and all the inputs u . This is so that we can eventually convert this function for $x_d[i]$ into a function for $x(t)$.

Unroll the implicit recursion you derived in the previous part to write $x_d[i+1]$ as a sum that involves $x_d[0]$ and the $u[j]$ for $j = 0, 1, \dots, i$.

For this part, feel free to just consider the discrete-time system in a simpler form

$$x_d[i+1] = ax_d[i] + bu[i] \quad (4)$$

and you don't need to worry about what a and b actually are in terms of λ and Δ .

(Hint: What is $x_d[1]$ in terms of $x_d[0]$? What is $x_d[2]$ in terms of (only) $x_d[0]$? What about $x_d[3]$? Can you find a pattern?)

From previous part: $x_d[i+1] = x_d[i] \cdot \underbrace{e^{\lambda\Delta}}_a + u[i] \cdot \underbrace{\frac{(e^{\lambda\Delta} - 1)}{\lambda}}_b$

$$\begin{aligned} x_d[1] &= a \cdot x_d[0] + b \cdot u[0] \\ x_d[2] &= a \cdot x_d[1] + b \cdot u[1] = a \cdot (a \cdot x_d[0] + b \cdot u[0]) + b \cdot u[1] \\ &= a^2 \cdot x_d[0] + u[0] \cdot ab + u[1] \cdot b \\ x_d[3] &= a \cdot x_d[2] + b \cdot u[2] \\ &= a \cdot (a^2 \cdot x_d[0] + u[0] \cdot ab + u[1] \cdot b) + b \cdot u[2] \\ &= a^3 \cdot x_d[0] + \underbrace{u[0] \cdot a^2 b}_1 + \underbrace{u[1] \cdot ab}_1 + u[2] \cdot b \end{aligned}$$

Using a summation: $x_d[i] = a^i \cdot x_d[0] + \sum_{j=0}^{i-1} u[j] \cdot a^{i-1-j} \cdot b$

- (c) For a given time t in continuous real time, what is the discrete i interval that corresponds to it?

(Hint: $\lfloor x \rfloor$ is the largest integer smaller than x .)

$$t = 3.5 \cdot 20 = 70$$

$$i = ? = \left\lfloor \frac{t}{\Delta} \right\rfloor = \left\lfloor \frac{70}{20} \right\rfloor = 3$$


- (d) Here's the first payoff! Use the results of part (a) and (b) to give an approximate expression for $x(t)$ for any t , in terms of $x_d[0] = x(0)$ and the inputs $u[j]$. You can assume that Δ is small enough that $x(t)$ does not change too much (is approximately constant) over an interval of length Δ .

(Hint: The assumption we just made allows us to approximate $x(t) \approx x\left(\Delta \left\lfloor \frac{t}{\Delta} \right\rfloor\right) = x_d\left[\left\lfloor \frac{t}{\Delta} \right\rfloor\right]$.)

↳ from $i = \left\lfloor \frac{t}{\Delta} \right\rfloor$ (c)

↳ from part (b): $x_d[i] = a^i x_d[0] + b \cdot \left(\sum_{j=0}^{i-1} a^{i-j-1} \cdot u[j] \right)$ (b) $a = e^{\lambda \Delta}$
 $b = (e^{\lambda \Delta} - 1) / \lambda$
 $x(t) \approx x\left[\Delta \cdot \left\lfloor \frac{t}{\Delta} \right\rfloor\right]$
 $= x_d\left[\left\lfloor \frac{t}{\Delta} \right\rfloor\right] = a^{\left\lfloor \frac{t}{\Delta} \right\rfloor} \cdot x_d[0] + b \cdot \left(\sum_{j=0}^{\left\lfloor \frac{t}{\Delta} \right\rfloor - 1} a^{\left\lfloor \frac{t}{\Delta} \right\rfloor - j - 1} \cdot u[j] \right)$

$$\Rightarrow x(t) \approx x_d\left[\left\lfloor \frac{t}{\Delta} \right\rfloor\right] = (e^{\lambda \Delta})^{\left\lfloor \frac{t}{\Delta} \right\rfloor} \cdot x_d[0] + \frac{(e^{\lambda \Delta} - 1)}{\lambda} \cdot \left(\sum_{j=0}^{\left\lfloor \frac{t}{\Delta} \right\rfloor - 1} (e^{\lambda \Delta})^{\left\lfloor \frac{t}{\Delta} \right\rfloor - j - 1} \cdot u[j] \right)$$



- (e) Now, we are going to turn this around. Suppose that the $u[i]$ is actually a sample of a desired input $u_c(t)$ in continuous time. Namely, suppose that $u[i] = u_c(i\Delta)$.

To clarify, $u(t)$ is a piecewise constant function; $u[i]$ is the discrete input that constructs $u(t)$; and $u_c(t)$ is the underlying input $u[i]$ is sampled from. **Solution:** In the terms of the earlier analogy, suppose $u_c(t)$ is a voltage signal (i.e. from another circuit) being read by the computer. At time $t = i\Delta$ the computer reads the input $u_c(i\Delta) = u[i]$ and provides the piecewise constant input $u(t)$ to the original circuit like before.

Notation
 $u_c(t)$: underlying signal
 $u(t)$: sampled
 $t \in [i\Delta, (i+1)\Delta)$
 $u[i]$

The underlying goal is to find an expression for $x(t)$ in the limit $\Delta \rightarrow 0$, in terms of $u_c(t)$ and the initial condition $x(0)$. To this end, start by substituting an appropriate value of u_c for u in the result from part (d). (Note: don't take any limits in this problem; just do the substitution.)

$$\Rightarrow x(t) \approx x_d\left[\left\lfloor \frac{t}{\Delta} \right\rfloor\right] = (e^{\lambda \Delta})^{\left\lfloor \frac{t}{\Delta} \right\rfloor} \cdot x_d[0] + \frac{(e^{\lambda \Delta} - 1)}{\lambda} \cdot \left(\sum_{j=0}^{\left\lfloor \frac{t}{\Delta} \right\rfloor - 1} (e^{\lambda \Delta})^{\left\lfloor \frac{t}{\Delta} \right\rfloor - j - 1} \cdot u[j] \right)$$

Substitute $u[j] \rightarrow u_c(t = j \cdot \Delta)$

$$x(t) \approx x_d\left[\left\lfloor \frac{t}{\Delta} \right\rfloor\right] = (e^{\lambda \Delta})^{\left\lfloor \frac{t}{\Delta} \right\rfloor} \cdot x_d[0] + \frac{(e^{\lambda \Delta} - 1)}{\lambda} \cdot \left(\sum_{j=0}^{\left\lfloor \frac{t}{\Delta} \right\rfloor - 1} (e^{\lambda \Delta})^{\left\lfloor \frac{t}{\Delta} \right\rfloor - j - 1} \cdot u_c(j \Delta) \right)$$

- (f) We want to take the limit $\Delta \rightarrow 0$ of our (discrete-time) expression and thus get a continuous-time function, but right now our discrete-time expression itself is pretty complicated. Let's simplify it by making some approximations which become exact in the limit.

Further approximate the previous expression by considering the following two estimates:

- Let $n = \lfloor \frac{t}{\Delta} \rfloor \approx \frac{t}{\Delta}$ where needed and treat $\Delta \approx \frac{t}{n}$. This is a meaningful approximation when we think about n large enough.
- Treat $\frac{1-e^{-\lambda\Delta}}{\lambda} \approx \Delta$. This is a meaningful approximation when we think about Δ small enough. One can derive this estimate by using Taylor's theorem from calculus, but it's not required here.

(Hint: Use the first estimate to get rid of "floor" terms, then use both estimates to simplify further.)

From part (e): $n = \lfloor \frac{t}{\Delta} \rfloor \approx \frac{t}{\Delta}$

$$\begin{aligned}
 x(t) &\approx x_d \left[\left\lfloor \frac{t}{\Delta} \right\rfloor \right] = (e^{\lambda\Delta})^{\left\lfloor \frac{t}{\Delta} \right\rfloor} \cdot x_d[0] + \frac{(e^{\lambda\Delta} - 1)}{\lambda} \cdot \left(\sum_{j=0}^{\left\lfloor \frac{t}{\Delta} \right\rfloor - 1} (e^{\lambda\Delta})^{\left\lfloor \frac{t}{\Delta} \right\rfloor - j - 1} \cdot u_c(j\Delta) \right) \\
 &\approx e^{\left(\frac{t}{\Delta} \cdot \frac{t}{\Delta}\right)} \cdot x_d[0] + \frac{(e^{\lambda\Delta} - 1)}{\lambda} \cdot \sum_{j=0}^{n-1} e^{\lambda\Delta \cdot \left(\frac{t}{\Delta} - j - 1\right)} \cdot u_c(j\Delta) \\
 &\approx e^{\lambda t} \cdot x_d[0] + \frac{(e^{\lambda\Delta} - 1)}{\lambda} \cdot \sum_{j=0}^{n-1} e^{\lambda t - \lambda\Delta j - \lambda\Delta} \cdot u_c(j\Delta) \\
 &= e^{\lambda t} \cdot x_d[0] + \frac{(e^{\lambda\Delta} - 1)}{\lambda} \cdot e^{\lambda t - \lambda\Delta} \cdot \sum e^{-\lambda\Delta j} u_c(j\Delta) \\
 &= e^{\lambda t} \cdot x_d[0] + \frac{1}{\lambda} \cdot e^{\lambda t} \cdot (e^0 - e^{-\lambda\Delta}) \\
 &\quad \hookrightarrow e^{\lambda t} \cdot \underbrace{\frac{1 - e^{-\lambda\Delta}}{\lambda}}
 \end{aligned}$$

$$x(t) = e^{\lambda t} x_d[0] + e^{\lambda t} \Delta \cdot \sum e^{-\lambda\Delta j} u_c(j\Delta)$$

in next few subproblems...

$$x(t) = e^{\lambda t} x_d[0] + e^{\lambda t} \int_0^t e^{-\lambda\tau} u_c(\tau) d\tau$$

(g) Here's our second payoff! We now obtain a continuous-time expression for $x(t)$, completing the transition into continuous-time. Take the limit of $x(t)$ as $\Delta \rightarrow 0$ or equivalently as $n \rightarrow \infty$. What is the expression you get for $x(t)$?

(*Hint*: Remember your definition of definite integrals as limits of Riemann sums in calculus.)