
EECS 16B

Spring 2022

Lecture 17

3/15/2022 

LECTURE 17 : - upper triangularization

Recall: diagonalization of an $n \times n$ matrix requires

n linearly independent vectors: $A\vec{v}_i = \lambda_i \vec{v}_i \quad i=1, \dots, n$

$$A \underbrace{[\vec{v}_1 \dots \vec{v}_n]}_{=: V} = [\lambda_1 \vec{v}_1 \dots \lambda_n \vec{v}_n] = \underbrace{[\vec{v}_1 \dots \vec{v}_n]}_{=V} \begin{bmatrix} \lambda_1 & & \\ & \ddots & \\ & & \lambda_n \end{bmatrix}$$

$$\Rightarrow V^{-1} A V = \begin{bmatrix} \lambda_1 & & \\ & \ddots & \\ & & \lambda_n \end{bmatrix}$$

Today: we can upper-triangularize even if we can't diagonalize. Upper-triangular form has some of the benefits of diagonal matrices:

1) Eigenvalues are the diagonal entries:

$$A = \begin{bmatrix} a_{11} & a_{12} & \dots \\ 0 & a_{22} & \\ \vdots & & \ddots \\ 0 & \dots & 0 & a_{nn} \end{bmatrix}$$

$$\lambda I - A = \begin{bmatrix} \lambda - a_{11} & -a_{12} & \dots & \\ 0 & \lambda - a_{22} & & \\ \vdots & & \ddots & \\ 0 & \dots & 0 & \lambda - a_{nn} \end{bmatrix}$$

$\lambda = a_{nn}$:
bottom
row is
zero;
row space
has $\dim < n \Rightarrow$ rank drops $\Rightarrow a_{nn}$ is
eigenvalue

$\uparrow$
 $\lambda = a_{11}$ zeroes out this
column $\Rightarrow$ rank drops
therefore a_{11} is an eigenvalue

If $\lambda = a_{ii}$
 $1 \leq i \leq n$

$$\lambda I - A = \begin{bmatrix} a_{11} - a_{11} & * & * & * \\ 0 & a_{22} - a_{22} & * & * \\ \vdots & \vdots & \vdots & \vdots \\ 0 & - & - & a_{ii} - a_{ii} & * \\ \vdots & \vdots & \vdots & \vdots & \vdots \\ 0 & - & - & - & a_{nn} - a_{nn} & * \end{bmatrix}$$

i columns;
 only top $i-1$ entries can be non zero
 $\Rightarrow$ can span at most the $i-1$ dimensional space
 spanned by first $i-1$ unit vectors
 $\Rightarrow$ these i columns are linearly dependent

$\Rightarrow$ Columns of $\lambda I - A$ are linearly dependent

$\Rightarrow$ not full rank $\Rightarrow \lambda = a_{ii}$ is an eigenvalue

2) Sol'n of vector diff. eq. or difference eq. broken down into scalar equations:

$$\frac{d}{dt} \vec{x}(t) = \begin{bmatrix} a_{11} & * & * \\ 0 & \ddots & * \\ \vdots & \ddots & \ddots \\ 0 & & 0 & a_{nn} \end{bmatrix} \vec{x}(t) + B \vec{u}(t)$$

$$\frac{d}{dt} x_n(t) = a_{nn} x_n(t) + (B \vec{u}(t))_n$$

$$x_n(t) = e^{a_{nn}(t-t_0)} x_n(t_0) + \text{term due to } (B \vec{u}(t))_n$$


$$\frac{d}{dt} x_{n-1}(t) = a_{n-1,n-1} x_{n-1}(t) + a_{n-1,n} x_n(t) + (B \vec{u}(t))_{n-1}$$

known function of time:
 treat as input to scalar diff. eq. for x_{n-1}

$$x_{n-1}(t) = e^{a_{n-1,n-1}(t-t_0)} x_{n-1}(t_0) + \underbrace{\int_{t_0}^t \dots}_{\text{Combined input terms from above go here}}$$

$$\frac{d}{dt} x_{n-2}(t) = a_{n-2,n-2} x_{n-2}(t) + a_{n-2,n-1} x_{n-1}(t) + a_{n-2,n} x_n(t) + \underbrace{(B \vec{u}(t))_{n-2}}_{\text{Combined input terms}}$$

Example: longitudinal motion of a car described by:



$$\frac{d}{dt} p(t) = v(t)$$

p : position
 v : velocity
 u : torque

$$M \frac{d}{dt} v(t) = \frac{1}{R} u(t)$$

$$\underbrace{\frac{d}{dt} \begin{bmatrix} p(t) \\ v(t) \end{bmatrix}}_{\vec{x}(t)} = \underbrace{\begin{bmatrix} 0 & 1 \\ 0 & 0 \end{bmatrix}}_{A_c} \underbrace{\begin{bmatrix} p(t) \\ v(t) \end{bmatrix}}_{\vec{x}(t)} + \underbrace{\begin{bmatrix} 0 \\ \frac{1}{RM} \end{bmatrix}}_{B_c} u(t)$$

Evalues of A_c : $\{0, 0\}$

Evector: $A_c \vec{v} = 0 \quad \vec{v} = \alpha \begin{bmatrix} 1 \\ 0 \end{bmatrix} \quad \alpha \neq 0$


no two lin. ind. evecs exist
 $\Rightarrow$ not diagonalizable
but in upper triangular form

Suppose $u(t) = \bar{u} = \text{constant}$. Solution of diff. eq.?

$$\frac{d}{dt} v(t) = \frac{1}{RM} \bar{u}$$

$$v(t) = v(t_0) + \frac{\bar{u}}{RM} (t - t_0)$$

$$\frac{d}{dt} p(t) = v(t) = v(t_0) + \frac{\bar{u}}{RM} (t - t_0)$$



$$p(t) = p(t_0) + v(t_0)(t - t_0) + \frac{1}{2} \frac{\bar{u}}{RM} (t - t_0)^2$$

Can find discrete time model if we set

$$t_0 = i\Delta, t = (i+1)\Delta, \bar{u} = u_d[i] :$$

$$\underbrace{p((i+1)\Delta)}_{p_d[i+1]} = \underbrace{p(i\Delta)}_{p_d[i]} + \underbrace{v(i\Delta)}_{v_d[i]} \Delta + \frac{\Delta^2}{2RM} u_d[i]$$

$$\underbrace{v((i+1)\Delta)}_{v_d[i+1]} = \underbrace{v(i\Delta)}_{v_d[i]} + \frac{\Delta}{RM} u_d[i]$$

$$\begin{bmatrix} p_d[i+1] \\ v_d[i+1] \end{bmatrix} = \underbrace{\begin{bmatrix} 1 & \Delta \\ 0 & 1 \end{bmatrix}}_{A_d} \begin{bmatrix} p_d[i] \\ v_d[i] \end{bmatrix} + \underbrace{\begin{bmatrix} \frac{\Delta^2}{2RM} \\ \frac{\Delta}{RM} \end{bmatrix}}_{B_d} u_d[i]$$

Next: prove any square matrix can be brought to an upper triangular form. Will prove for real matrices with real eigenvalues, but a complex version can be proven along the same lines.

Theorem: For any matrix $A \in \mathbb{R}^{n \times n}$ with real eigenvalues there exists orthogonal matrix $U \in \mathbb{R}^{n \times n}$ such that

$$U^{-1}AU = U^T A U$$

is upper triangular.

Thus, if $\frac{d}{dt} \vec{x}(t) = A \vec{x}(t) + B \vec{u}(t)$, then $\vec{y} = U^T \vec{x}$ satisfies

$$\begin{aligned} \frac{d}{dt} \vec{y}(t) &= U^T A \vec{x}(t) + U^T B \vec{u}(t) \\ &= \underbrace{U^T A U}_{\text{upper triangular}} \vec{y}(t) + U^T B \vec{u}(t) \end{aligned}$$

$\vec{x} = U \vec{y}$

We will prove the theorem by induction.

Proof by induction: Suppose we have a statement that depends on an integer $n=1, 2, 3, \dots, S_n$. To prove it by induction, show:

- S_1 is true
- for any $k \geq 1$, if we assume S_k is true then S_{k+1} is true.

Back to the Theorem: take the statement as S_n .

- Show S_1 is true: true b/c scalars are "upper triangular"
- Show if S_k is true, S_{k+1} is also true.

i.e.,

Assume: any real $k \times k$ matrix with real eigenvalues can be upper triangularized with orthogonal U .

Show: same is true for $(k+1) \times (k+1)$.

1. Let $A \in \mathbb{R}^{(k+1) \times (k+1)}$ be a matrix with real eigenvalues and let $\lambda_1, \vec{q}_1$ be an eigenvalue/eigenvector pair for A .

We will assume $\|\vec{q}_1\| = 1$. (No loss of generality because we can normalize the vector if its length is not 1 and it remains an eigenvector.)

2. Choose an orthonormal basis for $\mathbb{R}^{k+1}$ that includes $\vec{q}_1$: $\{\vec{q}_1, \vec{q}_2, \dots, \vec{q}_{k+1}\}$.

How to find them? Pick any k vectors in $\mathbb{R}^{k+1}$ which, when combined with $\vec{q}_1$, form a basis for $\mathbb{R}^{k+1}$. Then apply Gram-Schmidt with $\vec{q}_1$ as the first vector.

3. Then $Q = [\vec{q}_1 \dots \vec{q}_{k+1}]$ is an orthogonal matrix and

$$AQ = [A\vec{q}_1 \ A\vec{q}_2 \ \dots \ A\vec{q}_{k+1}]$$

$$= [\lambda_1 \vec{q}_1, A\vec{q}_2, \dots, A\vec{q}_{k+1}]$$

$$Q^T A Q = \begin{bmatrix} \vec{q}_1^T \\ \vec{q}_2^T \\ \vdots \\ \vec{q}_{k+1}^T \end{bmatrix} [\lambda_1 \vec{q}_1 \ A\vec{q}_2 \ \dots \ A\vec{q}_{k+1}]$$

$$= \begin{bmatrix} \lambda_1 \vec{q}_1^T \vec{q}_1 & * & \dots & * \\ \lambda_1 \vec{q}_2^T \vec{q}_1 & & & \\ \vdots & & & \\ \lambda_1 \vec{q}_{k+1}^T \vec{q}_1 & * & \dots & * \end{bmatrix}$$

by orthonormality

$$= \begin{bmatrix} \lambda_1 & * & \dots & * \\ 0 & * & \dots & * \\ \vdots & \vdots & & \vdots \\ 0 & * & \dots & * \end{bmatrix} \rightarrow \text{call this } \vec{p}^T$$

call this A_0 ($k \times k$)

$$Q^T A Q = \begin{bmatrix} \lambda_1 & \vec{p}^T \\ \vec{0} & A_0 \end{bmatrix} \quad \dots (1)$$

A_0 is not necessarily upper triangular but it is $k \times k$ so, by assumption at the top of last page, $U_0^T A_0 U_0$ upper triangular for some orthogonal U_0 .

(Note: we also need to know A_0 has real evlues.
That follows because $Q^T A Q$ has the same evlues as A
and because evlues of (1) are those of A_0 combined
with λ_1 .)

Define $U = Q \begin{bmatrix} 1 & 0 \\ 0 & U_0 \end{bmatrix}$ which is orthogonal:

$$\begin{aligned} U^T &= \begin{bmatrix} 1 & 0 \\ 0 & U_0^T \end{bmatrix} Q^T \Rightarrow U^T U = \begin{bmatrix} 1 & 0 \\ 0 & U_0^T \end{bmatrix} \underbrace{Q^T Q}_{=I} \begin{bmatrix} 1 & 0 \\ 0 & U_0 \end{bmatrix} \\ &= \begin{bmatrix} 1 & 0 \\ 0 & \underbrace{U_0^T U_0}_{I_{k \times k}} \end{bmatrix} = I_{(k+1) \times (k+1)}. \end{aligned}$$

$$\text{Note } U^T A U = \begin{bmatrix} 1 & 0 \\ 0 & U_0^T \end{bmatrix} \underbrace{Q^T A Q}_{\begin{bmatrix} \lambda_1 & \vec{p}^T \\ 0 & A_0 \end{bmatrix}} \begin{bmatrix} 1 & 0 \\ 0 & U_0 \end{bmatrix}$$

$$= \begin{bmatrix} \lambda_1 & \vec{p}^T \\ 0 & U_0^T A_0 U_0 \end{bmatrix} \begin{bmatrix} 1 & 0 \\ 0 & U_0 \end{bmatrix} = \begin{bmatrix} \lambda_1 & \vec{p}^T U_0 \\ 0 & \underbrace{U_0^T A_0 U_0}_{\substack{\text{upper triangular} \\ \text{also upper triangular}}} \end{bmatrix}$$

Summary: assuming that a $k \times k$ matrix

can be upper-triangularized, we have shown

the same is true for a $(k+1) \times (k+1)$ matrix.